absorption measurements were made, although exposures were in the low milliwatt/square centimeter range because they considered these levels to be nonthermalizing.

Contrary to the lack of engineering and physical scientific expertise associated with their microwave research, the biologists appeared capable of performing the required research. The major constraint in their biological capability seems not to be due to the quality of scientists but to the lack of up-to-date analytical equipment and instrumentation. Many of their laboratories are very limited in the amount of equipment, and much of what they have is obviously very old.

Recent visits to the Soviet Union have shown significant improvement in their exposure systems. Much of the improvement is probably due to recent visits to the United States and to observation of exposure systems in this country. The two institutes discussed above now have exposure rooms with absorbers on all walls and, in some cases, where multiple animals are being exposed, the exposure is from above. They also realize that much of their earlier electroencephalographic data, taken with metal implanted electrodes, are not valid because of potential artifacts produced by the interaction of the electrodes with the microwave fields.

During visits to the Soviet Union, some insight into the approach of the Soviet health institutes to studying this problem has been obtained. They believe that the more complex the biological system the greater the possibility of it being affected at low levels of exposure. Therefore, the health institutes in most cases use whole animal systems in their studies. They believe it is more important to obtain effects on different species of animals to extrapolate results from animal to man rather than to use *in vitro* or more simple systems to obtain basic mechanism of interaction data to enable such an extrapolation.

Adaptation is also an important consideration in their research. They state that human beings evolved under certain environmental conditions. Changes in these environmental conditions, increase or decrease, affect human beings, but human beings have adaptive capabilities which enables them to adapt to certain levels of changes without harmful effects. Levels greater in magnitude than those adaptive levels are defined as hazardous. Some of their research reports a biphasic change in a given biological parameter with increasing exposure time (see figure), which they explain by suggesting that the animal is attempting to adapt to microwave stress and overcompensates to the point that the change reverses direction (curve


ere in the low these levels to

ntific expertise peared capable n their biologibut to the lack Many of their , and much of

it improvement probably due to sure systems in exposure rooms iple animals are ze that much of netal implanted roduced by the

approach of the obtained. They reater the possiefore, the health ir studies. They erent species of an to use *in vitro* teraction data to

research. They ental conditions. decrease, affect es which enables il effects. Levels ed as hazardous. given biological iich they explain microwave stress direction (curve



Two suggested adaptation processes in response to microwave exposure.

1). Another form of adaptation which they discuss involves the idea that in the initial stages of exposure, animals can compensate for the microwave stress without changes in biological indicators. After a period of time, the animals can no longer adapt, and biological changes are then observed (curve 2).

Most Soviet research is performed, therefore, at low levels of exposure for long exposure duration. They consider that power densities above 1 mW/cm.<sup>2</sup> are high enough to produce harmful effects, and see no reason to perform research above this level. When we respond that in some of our experiments we see no significant changes at exposure levels above 1 mW/cm., their answer is that we did not expose long enough. Most of their low level experiments are for six months to a year, while American experiments are usually in terms of a few weeks. Long-term experiments using low levels of microwaves must be performed in the United States before the Soviet results can be verified or refuted.

### SOME REPORTED BIOLOGICAL EFFECTS

Before reporting on Soviet and other Eastern European biological effects, the results of the U.S.-U.S.S.R. cooperative program should be

briefly reviewed. In the early stages of the cooperative program, it consisted mainly of an exchange of results on projects related to the central nervous system and behavior. American research consisted primarily of acute experiments with exposure levels generally of 5 mW/cm.<sup>2</sup> and above, while Soviet experiments were long-term, low-level experiments at 500 μW/cm.<sup>2</sup> and below. At the end of the first year of the cooperation, the Soviets reported changes in bioelectric brain activity at 10, 50, and 500  $\mu$ W/cm.<sup>2</sup> in rats and rabbits exposed for 7 hours/day for 30 days to 2,375 MHz. radiation. Levels of 10 and 50  $\mu$ W/cm.<sup>2</sup> stimulated brain activity, while 500  $\mu$ W/cm.<sup>2</sup> suppressed activity as seen from an increase of slow, high amplitude  $\Delta$ -wave in rabbits. At 500  $\mu$ W/cm.<sup>2</sup> a decrease in capacity for work, in investigative activity, and sensitivity to electric shock threshold in rats were reported. Research by American investigators on rats exposed to 5 mW/cm.<sup>2</sup> for shorter durations of exposure to 2,450 MHz. radiation showed no statistical difference in electroencephalogram, no change in locomotive activity in a residential maze, and no change in performance on a fixed ratio schedule of reinforcement below 5 mW/cm.<sup>2</sup> (0.5 and 1.0 mW/cm.2) but a trend toward decrease in performance at 5 mW/cm.<sup>2</sup> and a large decrease in performance at 10 and 20 mW/cm.<sup>2</sup>

It became obvious that, except for our being more familiar with their experimental design, we were no closer to understanding differences between American and Soviet results. It was then decided to perform a duplicate experiment to determine whether similar effects could be observed. Rats were exposed from above for seven hours/day, seven days/ week for three months to 500  $\mu$ W/cm.<sup>2</sup>. Dr. Richard Lovely of the University of Washington, project leader on the duplicate project, spent four weeks in the Soviet Union to observe the behavioral and biochemical tests performed on the animals. The American study found a drop in sulfhydryl activity and blood cholinesterase as reported in the Soviet study. Blood chemical analyses at the termination of three months exposure indicated that exposed animals, relative to controls, suffered from aldosteronism. The latter interpretation of the high sodium-low potassium levels found in the blood was confirmed by necropsy and histopathologic study of the adrenal glands, revealing that the zona glomerulosa was vacuolated and hypertrophied. In addition, all behavioral parameters assessed at the end of three-month exposures revealed significant differences between groups in the same direction as those reported in the Soviet study, i.e., increased threshold to footshock detection, decreased activity in an

ogram, it conto the central d primarily of m.2 and above, riments at 500 ooperation, the 0, 50, and 500 ) days to 2,375 brain activity, crease of slow, ease in capacity electric shock investigators on osure to 2,450 encephalogram, id no change in low 5 mW/cm.2 erformance at 5 ) mW/cm.<sup>2</sup>

miliar with their differences beed to perform a its could be obday, seven days/ i Lovely of the te project, spent and biochemical found a drop in ed in the Soviet ree months expois, suffered from ım-low potassium id histopathologic glomerulosa was al parameters asificant differences the Soviet study, sed activity in an

TABLE I. FREQUENCY OF HYPOTONIA IN PERSONNEL EXPOSED TO MICROWAVES ACCORDING TO SOVIET AUTHORS

Author and year	No. of individuals examined	Frequency of hypotonia %
Kerovski, A.A., 1948	87	38
Osipov, Ju A., 1952	108	30
Shipkova, V.A., 1959	110	20
Orlova, A.A., 1960	525	26-33, various groups
Uspenskaja, I.V., 1961	100	30
Volfavskaja, R.N. et al., 1961	101	27-45,
		various groups
Frelova, L.T., 1963	172	25.6
Komarov, F.I. et al., 1963	53	22.6
Gembricki, E.V., 1966	210	14

Reproduced by permission from Baranski, S. and Czerski, P.: Biological Effects of Microwaves. Stroudsburg, Pa., Dowden, Hutchinson and Ross, 1976.

open field, and poorer retention of an avoidance response when reassessed following conditioning. This replication of the Soviet results at 500  $\mu$ W/cm.<sup>2</sup> emphasizes the need for additional long-term, low-level microwave bioeffects research.

Effects on human beings. Most of the bioeffects research in Eastern European countries is performed by Soviet and Polish investigators. A small amount of data, mainly effects on human workers, have been reported by Czechoslovakian scientists. Reported effects on personnel occupationally exposed to microwaves has been summarized by Baranski and Czerski<sup>2</sup> and are shown in Tables I and II. These results come primarily from two groups in the Soviet Union who have done extensive clinical studies on microwave workers. Baranski and Czerski<sup>2</sup> refer to the groups as the Moscow group and the Leningrad group. The Moscow group has reported changes in white blood cells, although no characteristic changes in peripheral blood have been found. Blood proteins, serum histamine content, and enzyme activity have been reported but they returned to normal levels after a few days of rest. The influence of microwaves on the autonomic nervous system was evidenced by reports of vagotonia, bradycardia, and hypotonia. With periodic exposure to power density levels of 0.1 to 10 mW/cm.<sup>2</sup>, marked changes in cardiac rhythm (variability or pronounced bradycardia) were reported. Hypotonia was one of the more pronounced effects (Table I). Prolonged exposure to power density levels from 0.01 to 0.1 mW/cm.2 produced effects similar to those

Table II. COMPLAINTS OF MICROWAVE WORKERS (%) ACCORDING TO VARIOUS AUTHORS

Author and year	Headaches	Fatigue dispro- portional to effort	Sleep distur- bances	Irritability	Abnormal sweating	No. of workers examined
Uspenskaja, N.V., 2963	37	31	29	9	7	100
Sadtchikova, M.N., 1963 (various groups)	12-39	20-35	_	8-27	_	447
Kilmkova-Deutschova,	43	39	35	<del></del>		73
1963 (Czech.)	43	4	45	10		103
Serel, 1959 (U.S.S.R.) Fjagin, N.V., 1966 (U.S.S.R.)	33.5	46.2	25.3	9.6	25.5	573
Ramzen-Evdokimov and Sorokin, 1970 (U.S.S.R.)	44	29	35	36	25	155
Controls	1.5	22	2	10		100
Uspenskaja, 1963	15 8		2	8		100
Sadtchikova, 1963 Tjagin, 1966	8 10.8	10 5.9	8.7	<del></del>	2.7	184
Ramzen-Evdokimov and Sorokin, 1970	7	8	3	<del></del>	4	50

Reproduced by permission from Baranski, S. and Czerski, P.: Biological Effects of Microwaves. Stroudsburg, Pa., Dowden, Hutchinson and Ross, 1976.

mentioned above. The symptoms were less pronounced and reversible after periods of rest.

The Moscow group also reports that chronic exposure leads to a neurocirculatory syndrome in three stages. The first symptoms are reversible and include changes in blood pressure and cardiac rhythm. In the second stage, cardiovascular changes become more pronounced, and electroencephalographic changes which indicate disturbance of the diencephalan were also found. Hyperactivity of the thyroid is also indicated. A condition defined in Soviet literature as an "asthenic state" may develop. Such subjective complaints as headaches, excitability and irritability, fatigue, and pains around the cardiac region (Table II) characterize this stage. The third stage of effects is characterized by a greater magnitude of change in the complaints and symptoms and additional electroencephalographic changes.

The Leningrad group discusses their findings in terms of an acute microwave syndrome and a chronic microwave syndrome. The acute syndrome is characterized by subjective complaints of headaches, nausea, vertigo, and sleep disturbance. Hypertonia, changes in cardiac rhythm, skin rash, and decrease in amplitude of electroencephalographic alpha waves are also characteristic. Symptoms are transient and disappear completely after rest. The chronic syndrome is expressed in terms of adaptive processes. Subjective complaints of headaches, fatigue, and the like occur during the first three months of exposure, and reappear about the sixth to eighth month. After the first year a period of adaptation occurs which varies in duration. The complaints and symptoms of neurovegetative disturbances reappear after five years of work (chronic overexposure syndrome). This chronic overexposure syndrome is characterized by complaints of headaches, irritability, sleep disturbances, weakness, decrease of sexual activity, pains in the chest, and a general unhealthy feeling ("a general feeling of ill-being").

A large number of Polish clinical studies confirm the Soviet findings on the periodicity of the subjective complaints for exposures to 1 to 10 mW/cm.<sup>2</sup>.<sup>2</sup> Subjective complaints were exactly the same as described by the Soviet authors (Table II), and were observed in up to 70% of persons exposed to the 1 to 10 mW/cm.<sup>2</sup> power density levels. The few Czechoslovakian reports (Klimkova-Deutschova<sup>1</sup> and Serel<sup>2</sup>) agree with the Soviet authors (Table II). They also discuss symptoms in terms of a "neurasthenic syndrome."

### TABLE III. SOME DATA ON THE EXPOSURE OF MAN AND ANIMALS TO MICROWAVE FREQUENCY FIELDS (ARRANGED ON INTENSITY SCALE)

Power density		
W/cm. <sup>2</sup>	ı	Eye cataracts in dogs after exposure for three to five hours
	800	
	600	(L,M) Pain sensation during exposure
	300	Brief increase in blood pressure; after 20 to 60 minutes marked decrease (cat, rabbit, dog)
	200	(L) Malformation of offspring after exposure for 10 to 15 minutes (chicken eggs, $\lambda = 12.6$ cm), death of cats and rabbits (t = 20 to 60 minutes). Reduction of redox in tissue
mW/cm.2		
	100	(M) Increase in blood pressure with subsequent marked decrease; in case of chronic exposure—stable hypoxia. Stable morphologic changes in the cardiovascular system. Bilateral cataracts.
	40	(L) Increase in blood pressure with subsequent marked decrease; multiple hemorrhages, $\lambda=3$ to 10 cm, in liver (dilatation of vessels and hemorrhages $\lambda=10$ cm.). Increase in blood pressure of 20 to 30 mm. Hg (exposure for 0.5 to 1.0 hour).
	10	(M) Changes in conditioned reflex activity, morphological changes in cerebral cortex (L). Vague shifts in blood pressure (exposure time 150 hours), change in blood coagulability. Hyperplasia of liver cells, $\lambda=3$ to 10 cm. (chronic exposure). ECG changes (wavelengths other than DTsV—expansion not given). Change in receptor apparatus.
	5	Threshold intensity at which there are changes in the testis and blood pressure changes (multiple exposure). Brief leukopenia and erythropenia. Darkening of the crystalline lens.
	3	(M) Decrease in blood pressure, tendency to quickening of pulse, fluctuation of cardiac blood volume.
mW/cm.2		
	1	(M) Decrease in blood pressure, tendency to quickening of pulse, insignificant variations in cardiac blood volume. Decrease in blood pressure level, decrease in ophthalmotone (t: daily, 3.5 months). Disadaptation, disorders of immunological protection control mechanisms (L).
	400	Depression of secretions in dogs
	300	(L,M) Some changes in the nervous system in case of exposure for 5 to 10 years
	200	Changes in function of neurons in dogs
	100	(L) Tendency to decrease in blood pressure with chronic exposure
	40	(L) Tendency to decrease in blood pressure with chronic exposure
$\mu$ W/cm. <sup>2</sup>		
	20	(M) Thinning of pulse, tendency to a decrease in arterial pressure. Cases of body sensitivity observed. Well-expressed increase in skin temperature in persons earlier exposed.
	M	Data applies to man-all other to animals

Source: Minin, B.A.: Microwaves and Human Safety. U.S. Joint Publication Service, JPRS 65506. August 20, 1975.

L Lowest power density indicated by authors

Vol. 55, No. 11, December 1979

pressure. ease in skin exposure

exposure

sposure for

SCALE)	LS TO

i minutes = 20 to 60

lecrease; phologic

### Table IV. SOME RESULTS OF EXPERIMENTAL STUDIES ON THE BIOLOGICAL EFFECTS OF VERY LOW INTENSITY MICROWAVES (UP TO 150 $\mu$ W/cm.<sup>2</sup>)

of pulse,

is and venia and

iposure asia of ages (wave-

Investigated function	Radiation intensity µW/cm.²	Character of changes	Investigator
Body weight	150	Lag in weight (chronic experiment)	V.V. Markov
Arterial pressure	150	Biphasic course with marked hypotension (chronic experiment)	V.V. Markov
Reproductive function	150	Decreased fertility, decreased litter size, increased number of defective progeny, increased embryonic mortality etc. (chronic experiment)	A.N. Bereznitskaya et al.
Central nervous system	10-20 and higher 150	<ol> <li>EEG changes with predominant synchronization (acute experiment)</li> <li>Bivariant shifts with predominance of activation (acute experiment)</li> <li>Bivariant shifts in the subcortical-basal structures (chronic experiment)</li> </ol>	Z.V. Gvozdikcova et al.
Electromyography	150	Increased electrical activity of active unit	V.V. Markov
Hypothalamus- drenal cortex System	150	<ol> <li>Weight change of endocrine glands (hypothysis adrenals)</li> <li>Change in the neurosecretory function of the hypothalamus.</li> <li>Tendency for increased levels of norepinephrine in the adrenals</li> </ol>	N.K. Demokidova
Metabolism	150	Changes in water and electrolyte metabolism (Na, K, water, and total nitrogen excretion)	N.K. Demokidova
Immunology	150	Inhibition of neutrophils phagocytic activity	A.P. Vokova and V.V. Markov

Source: Gordon, Z.V., editor: Biological Effects of Radiofrequency Electromagnetic Fields. U.S. Joint Publication Service, JPRS 65506, August 20, 1975.

TABLE V. RADIOFREQUENCY STANDARDS FOR OCCUPATIONAL EXPOSURE TO CONTINUOUS WAVE FIELDS (WORKING DAY)

Country	Radiation frequency (GHz.)	RFR intensity (mW/cm.²)
United States		
ANSI*	0.01-100 GHz.	10
OSHA†	0.01-100 GHz.	10
ACGIH‡	0.1 -100 GHz.	10
Great Britain	0.3 - 30 GHz.	10
Сапада	0.1 -100 GHz.	10
		(1 proposed)
Sweden	0.3 - 3  GHz	1 '
Poland	0.03-300 GHz.	0.2
Czechoslovakia	0.3 -300 GHz.	0.025
U.S.S.R.	0.3 -300 GHz.	0.01

<sup>\*</sup>American National Standards Institute

†Occupational Safety and Health Administration

‡American Congress of Governmental Industrial Hygienists

Effects on animals. A compilation of many Soviet reports on the effects of microwave radiation on animals is presented (Tables III and IV). Both Soviet and Polish investigators report behavioral, central nervous system, cardiovascular system, biochemical, endocrine and metabolic, hematological and reproductive functional changes in animals. All of these reported effects (Tables III and IV) were observed at power densities equal to or less than  $10~\text{mW/cm}.^2$  and many of the effects were observed at power densities less than  $1.0~\text{mW/cm}.^2$  Biological changes in both human beings and animals have been reported below  $500~\mu\text{W/cm}.^2$ 

### MICROWAVE EXPOSURE STANDARDS

The current standards for occupational exposure to continuous wave fields for a working day show large differences between Soviet and other Eastern European countries and western countries (Table V). Although Gordon<sup>3</sup> also presents a permissible exposure level of 1  $\mu$ W/cm.<sup>2</sup> for the general population as well as the 10  $\mu$ W/cm.<sup>2</sup> for occupational exposure, the Soviet Union does not have at this time a "legal" permissible general population exposure levels. However, the Institute of General and Communal Hygiene in Kiev, under the direction of the Soviet Ministry of Health, has proposed a general population standard for microwave exposure of 5  $\mu$ W/cm.<sup>2</sup> (Table VI). This maximum permissible level is enforced at present and will become an official standard at the end of this

TA

Ra

Sh-Ul Mı

y٠

ot m

Ci

SE

le

al

n

le

S

p:

p

st ti

iı

g

tÌ

tì

Note: Some countries have more permissive intensities for shorter exposure times and more restrictive intensities for exposure of the general population.

)NAL AY)

nsity

sed)

ore restrictive

he effects
V). Both
s system,
matologireported
ual to or
at power
in beings

us wave and other Although <sup>2</sup> for the xposure, general d Comnistry of re exposit is end of this

\cad. Med.

TABLE VI. MAXIMUM ALLOWABLE LEVELS (MAL) OF ELECTROMAGNETIC ENERGY IN THE HUMAN SETTLEMENTS

Radiowaves ranges	Limits of ranges (frequency, wavelength)	MAL of EMW energy on the dwelling territory
Long waves	30-300 kHz. (10-1 km.)	20 V/m.
Middle waves	0.3-3 MHz. (1-0.1 km.)	10V/m.
Short waves	3-30 MHz. (100-10 m.)	4 V/m.
Ultrashort waves	30-300 MHz. (10-1 m.)	2 V/m.
Microwaves 24-hour exposure	300 MHz300 GHz. (1 m1 mm.)	$5 \mu \text{W/cm.}^2$

year unless reasons not to accept it are presented to the Ministry of Health.

Differences in the meaning of standards exist between the Soviet and other Eastern European countries and the United States. First, the maximum permissible exposure level is set at the value where no biological effects occur. No differentiation between effects and hazards is made in setting their standards. Although most of their reported effects at low levels are reversible and return to normal levels after a period of time, and although they recognize that a difference between an effect and a hazard might exist, these facts are not considered in setting maximum permissible levels. Second, the occupational and general population standards in the Soviet Union and other Eastern European countries are not applicable to persons in military and space programs. This enables them to avoid the problems of restrictive use in high priority applications when setting very stringent maximum permissible exposure levels for lower priority situations. This exclusion of individuals in certain work activities is of course impossible in the United States if harmful effects do in fact occur at a given level and duration of exposure. Unfortunately, with the data base that exists today, this "safe" level cannot be set with certainty for either the general population or occupational groups.

#### SUMMARY

The Soviet and other Eastern European countries have published much information on the biological effects of microwave radiation. Almost no contact between Eastern European and Western scientists on this problem existed prior to 1972, when the cooperative agreement on environmental

health was signed with the Soviet Union. Since 1972 scientific exchanges have occurred on an annual basis with some American scientists staying as long as six weeks to observe Soviet research. Other exchange programs between Polish investigators and American agencies have taken place over the past several years. The exchanges have provided us with a better insight into research in these countries.

Because of my direct involvement with the U.S.-U.S.S.R. cooperative program, this paper has concentrated on Soviet research. These institutes stress human clinical studies and long-term, low-level effects on whole animal systems. They report an "asthenic syndrome" in occupationally exposed workers, characterized by headaches, irritability, fatigue, sleep disturbances, pains in the chest, and a general feeling of ill-being. Polish and Czechoslovakian investigators report the same human effects as the Soviet investigators. Reports of animal studies by Soviet and Polish investigators at exposure power densities of 10 mW/cm.<sup>2</sup> and below include effects on central nervous system, behavior, cardiovascular system, hematology and blood biochemistry, immunology, endocrinology and metabolism, and reproductive function.

Maximum permissible exposure levels in the Soviet Union and other Eastern European countries and accepted safe levels in the United States differ drastically. For a full working day of exposure, the Soviet standard is 1,000 times less than that presently accepted in the United States. However, the Soviet level is set where supposedly no biological changes occur, even if reversible. No evaluation of the biological changes in terms of harmful consequences is incorporated into the standard. It should also be noted that certain programs (military and space) do not adhere to the same requirements as those set by the Soviet Ministry of Health.

### **Ouestions and Answers**

DR. RUSSELL CARPENTER: Dr. McRee, were methods of measuring the microwave levels of power what you showed us, how can we really take any stock in what they offer? When they say "low levels" and one measures it with plaster walls and no anechoic chambers, I do not know what their device was. Do you think their measurements are very reliable?

DR. McRee: The PO-1 meter that they used is a reliable device for measuring incident power density. If they were exposing the animal at 1 or 10 mW/cm.<sup>2</sup> and one had a 10 to 100 amplification factor or error factor in

th

th

W

 $a^1$ 

11

re T

a

d

exchanges staying as programs place over a better

institutes on whole pationally ue, sleep g. Polish its as the sh inves-/ include system, and me-

nd other ed States standard States. changes in terms uld also e to the

ing the ly take nd one t know iable? ice for at 1 or ctor in

the measurement, obviously it could be a high level thermal effect. But they report biological changes at 10  $\mu$ W/cm.<sup>2</sup> and I just cannot see that much error. I would estimate a factor of 10 in that particular exposure. Dr. William Guy and I had some meters over there but we, unfortunately, were not allowed to take some measurements that we wanted, but one can look at the reflectivity and so forth at 10  $\mu$ W incident and see no greater increase than about 100 µW to 1 mW. They do not report specific absorption rates. In fact, I have never seen them measure or attempt to measure these. Because they say that at their levels they have no heating. I could not measure a specific absorption rate at 10  $\mu$ W/cm.<sup>2</sup> So they don't really feel there is a need. They have improved their exposure systems. The plaster walled room shown in the slide with the animal in front of the diathermy unit has now been replaced by a standard gain horn irradiator. animals in the far field, and microwave absorbers on the walls. We have done a duplicate project with them. At exposure to 500  $\mu$ W/cm.<sup>2</sup> for seven hours per day, seven days a week, for three months, we examined blood chemical and behavioral changes. Behavioral changes measured were almost identical to those measured by Shandala's group. This work at the University of Washington by Dr. Lovely, I think, amplifies the problem, or rather the realities, of the situation. We must do some chronic low level work, and we might be surprised. I think that Lovely and Guy were probably more surprised than anyone else that they found those results at a half mW/cm.<sup>2</sup>

MR. EDWARD L. HUNT (Walter Reed Army Institute of Research): The Russians have begun to adopt some of the same essential kinds of specifications that we use. For example, Tyazhelov in Helsinki has used watts per kilogram, which is our specific absorption rate (SAR).

DR. McRee: They have begun to do this, particularly where it is relevant, where they think they can measure specific absorption rates. Of course, one can measure such rates in any condition if one just exposes at a higher power density level and extrapolates down to the low level at which one tries to do the experiment. But one encounters heat transfer conduction problems because one is basically measuring the temperature.

MR. GEORGE WILKENING: Is it a valid argument not to use SAR because of heat? One doesn't necessarily have to have it, does one?

DR. McRee: One must measure the fields or the temperature and unless the BRH field probe is as good as they say, and is made available....

DR. JAMES FRAZER: Dr. McRee, you are only halfway to the solution.

When you measure SAR, you indeed measure the electric field in the tissues.

DR. McRee: I agree, but presently most people obtain SAR from temperature measurements. Most people do not have a field probe with which to measure local fields. Most do have a temperature measuring device.

DR. DON R. JUSTESEN: Does this not imply that human beings are more sensitive to weak electromagnetic fields than the best sensors that the engineers can devise? Presumably, it is beyond the capability of the engineers to detect these fields, while the human being not only reacts to them but suffers terribly from all manner of neurasthenic illness. Is this not the implication?

DR. McRee: You are trying to make a point. What is it?

DR. JUSTESEN: It is possible that the central nervous system is the most sensitive of all instruments to weak radiofrequency fields—that we could throw away our power-density meters and more accurately index the presence of extremely weak but nonetheless harmful fields simply by counting the number of factory workers or embassy employees with neurasthenic symptoms.

MR. WILKENING: To use the incidence of a malady in which the role of microwaves is at best conjectural as an index of field strength would certainly beg the causal question.

DR. JUSTESEN: Precisely my point. I spoke with tongue in cheek. Although I do not disparage the sensitivity of the nervous system to electromagnetic fields, I do disparage the tendency of some individuals to blame any manifestation of disease on the microwaves. A good case in point is Dr. Milton Zaret's assertion<sup>5</sup> that over-the-horizon radar beams from the Soviet Union are responsible for the high incidence of cardiac disease and heart failure in the North Karelia district of Finland. There is little doubt that the Soviet-launched radiation incident on the inhabitants of this district is unmeasureably small—much less than the microwave radiation emitted by the human body. One must disparage the assertion of an etiological role of any agent in any pathological condition in the absence of measurement, correlation, and verification.

DR. McRee: I understand the point Dr. Justesen makes, and it is a valid one. We should not jump to the conclusion that health effects found in persons who might be in a location where microwaves are present are due to the microwave radiation. It is important to characterize the microwave

radiatent. I low p at the experiment of the notate of the

1. Bic Mi tion Wa

2. Ba

Eff

Do

3. Go of Fig.

d in the

R from be with easuring

re more hat the of the eacts to this not

> ie most e could ex the ply by s with

> > role of would

cheek. em to ials to ase in eams ardiac ere is nts of adiaof an ce of

> valid id in due vave

radiation levels and any other environmental factors which might be present. However, the Soviets can measure accurately microwave levels in the low  $\mu$ W/cm.<sup>2</sup> range and have conducted long-term experiments on animals at these levels. Reversible biological changes have been reported in these experiments at power density levels as low as 5µW/cm.<sup>2</sup>

DR. JUSTESEN: Dr. Zorach Glaser, formerly with the Department of the Navy, and now in transition from the National Institute of Occupational Safety and Health to the FDA's Bureau of Radiological Health, is owed a debt of gratitude by all physicians, biologists, and biological engineers who labor in the radiofrequency field. For years Dr. Glaser has performed the nearly thankless task of cataloging, updating, and disseminating titles and sources of published information on biological effects of radiofrequency fields. His current compendium exceeds 5,000 titles. Someone should acknowledge Dr. Glaser's onerous but highly useful labor.

### REFERENCES

- 1. Biologic Effects and Health Hazards of Microwave Radiation. Proc. International Symposium, Warsaw, Poland. Warsaw, Polish Med. Pub., 1974.
- √2. Baranski, S. and Czerski, P.: Biological 5. Zaret, M. M.: Electronic smog as a poten-Effects of Microwaves. Stroudsburg, Pa., Dowden, Hutchinson and Ross, 1976.
- 3. Gordon, Z.V., editor: Biological Effects of Radiofrequency Electromagnetic Fields. U.S. Joint Publications Research
- Service, JPRS 63321, October 30, 1974.
- 4. Minin, B.A.: Microwaves and Human Safety, U.S. Joint Publication Service. JPRS 65506, August 20, 1975.
- tiating factor in cardiovascular disease: A hypothesis of microwaves as an etiology for sudden death from heart attack in North Karelia. Med. Res. Eng. 12:13-16, 1976.



### **INTRODUCTION**

Welcome to the 15th Annual International Symposium on Man and His Environment in Health and Disease. This Symposium is one of the most advanced forums in the world addressing the research and treatment of environmental effects on health and disease.

At this year's Conference, experts from throughout the world will share their extensive experience and specialized knowledge with an audience of physicians, scientists and health professionals.

### SPECIAL FOCUS

The 1997 Annual International Symposium will focus on the Environmental Aspects of EMF and Bioelectricity. This Conference will explore some of the latest data and provide a forum for discussion as well as case studies to help the professional.

### GOALS OF THIS SYMPOSIUM

- 1.) To provide the practicing physician with a clear definition of Bioelectricity and EMF.
- 2.) To improve the attendants' awareness of the effects of Bioelectricity and EMF on the human body.
- 3.) To develop EMF models for clinical use.
- 4.) Present treatment modalities for the sensitive patient.

### **OBJECTIVES OF THE SYMPOSIUM**

Upon completion of this conference, the physician should be able to:

- 1.) To provide an overview and a greater appreciation of EMF and Bioelectricity.
- 2.) Understand the problems associated with EMF and Bioelectricity.
- 3.) Be able to identify, diagnose, treat and prevent these environmentally triggered responses to the human body.

### **CONFERENCE FORMAT**

The AEHF Committee in conjunction with the AAEM has selected some of the leading experts in the field of EMF and Bioelectricity.

Each speaker's presentation will last approximately 20 minutes and will be followed by a 10 minute question and answer session. All speakers are encouraged to use any and all appropriate audio/visual aids. (A brief outline of the speech is included in this booklet.)

Every afternoon, we will have a case study/panel discussion. This session will consist of various faculty members discussing real cases. The audience is encouraged to participate in these discussions.

### **ACCREDITATION**

This CME activity has been planned and implemented in accordance with the Essentials and Standards of the Accreditation Council For Continuing Medical Education (ACCME) through the partnership of the American Academy of Environmental Medicine (AAEM) and the American Environmental Health Foundation.

The American Academy of Environmental Medicine is accredited by the Accreditation Council For Continuing Medical Education to provide CME activities for physicians.

The American Academy of Environmental Medicine designates this CME activity for 24 credit hours in Category I of the Physicians Recognition Award of the American Medical Association.

### COMMERCIAL SUPPORTERS OF THE ANNUAL INTERNATIONAL SYMPOSIUM

Thanks to all of the companies listed below that have supported our Annual International Symposium over the past years.

### 10 YEARS OR MORE OF CONTINUED SUPPORT

- ALLERGY RESEARCH GROUP
- ALLERMED CORPORATION
- ANTIBODIES ASSAY LABORATORIES
- DOCTOR'S DATA
- ENVIRONMENTAL HEALTH CENTER-DALLAS
- KLAIRE LABORATORIES

### **5 YEARS OR MORE OF CONTINUED SUPPORT**

- ABRAMS ROYAL PHARMACY
- ACCU-CHEM
- AFM ENTERPRISES
- AEROBIC ENTERPRISES
- BIO-TECH
- BIOTICS RESEARCH CORPORATION
- IVP PHARMACEUTICAL CARE, INC.
- PHYSICIAN SALES AND SERVICES
- R H OF TEXAS
- W.B. SAUNDERS
- TEXAS POWER VAC

### GIVEN IN COOPERATION WITH

William J. Rea, M.D., F.A.C.S.

Symposium Chairman

Environmental Health Center - Dallas

Dallas, Texas

Alfred R. Johnson, D.O.,
Gerald H. Ross, M.D.,
Bertie Griffiths, Ph.D.
Environmental Health Center - Dallas
Dallas, Texas

Ervin J. Fenyves, Ph.D.

Department of Physics and Environmental Sciences
University of Texas at Dallas
Richardson, Texas

Allan Lieberman, M.D.

The Center for Environmental Medicine
N. Charleston, South Carolina

Gary R. Oberg, M.D.
Chairman, AAEM CME Committee
Crystal Lake, Illinois

## THE FOLLOWING ORGANIZATIONS ARE SPONSORS FOR THE 15th ANNUAL SYMPOSIUM

### GOLD

ALLERMED CORPORATION

DOCTOR'S DATA

RED OAK DRUG

### SILVER

PHYSICIANS SALES AND SERVICES
R H of TEXAS

## **IMPLANTS**

A Public Conference

# Sponsored By: American Environmental Health Foundation

Guest Speakers:

Deborah Baird, M.D. - Pediatrician "TMJ IMPLANT SYNDROME; IT'S NOT ALL IN YOUR HEAD"

Arthur D. Ericsson, M.D. - Neurologist "SYNDROMES ASSOCIATED WITH SILICONE BREAST IMPLANTS"

James Key, M.D. - Orthopedic Surgeon "ORTHOPEDIC IMPLANTS"

William J. Rea, M.D. - Cardiovascular Surgeon "OVERVIEW OF THE IMPLANT SYNDROMES: BREAST, VASCULAR, ABDOMINAL, EYES, PACEMAKER, TREATMENT POST REMOVAL"

Sherry Rogers, M.D. "The Causes and Cures of Breast Cancer"

Larry Wolford, D.D.S.
"LOCAL AND SYSTEMIC EFFECTS OF PROPLAST / TEFLON IMPLANTS
TO THE TMJ"

+++

Saturday, February 22, 1997 The Le Meridien Hotel - Dallas (Cambridge Room)

### 1997 SYMPOSIUM FACULTY

Jacques Benveniste, M.D. I.N.S.E.R.M. U 200 32 Rue Des Carnets 92140 Clamart

Phone: Fax: 011 33 1 46 32 12 07

: 011 33 1 46 31 02 77

Joel R. Butler, Ph.D.

Phone:

918/534-0190

Environmental Health Psychologists

Fax:

918/534-1071

P.O. Box 399 Dewey, OK 74029

France

### General Information:

- Received doctorate in clinical psychology from Louisiana State University, Baton Rouge, Jan. 1962. Served as Professor LSU Baton Rouge; University of Georgia, Athens; Virginia Commonwealth University, Richmond, (Clinical Director); University of North Texas, Denton (Professor and Chair--Retired).

### Additional Information:

- Published/presented over 100 professional papers and contributor/author to 10 books.

Nancy A. Didriksen, Ph.D. Health Psychology Behavioral Medicine Associates 100 N. Cottonwood Dr., Suite 106 Richardson, TX 75080 Phone: Fax: 972/889-9933

972/889-9935

Dr. Didriksen received her degree in health Psychology/Behavioral Medicine from the University of North Texas in 1986. She completed her clinical internship in the Environmental Control Unit at Northeast Community Hospital in Bedford, Texas and has been evaluating and treating environmentally sensitive patients since that time.

She has given many presentations and has published on the psychological aspects of environmental illness and is the first author of a chapter titled <u>Psychological Concomitants of Chemical Sensitivity: Evaluation and Treatment</u> in Dr. William J. Rea's Vol. 4, <u>Chemical Sensitivity</u>. She is an adjunct professor of psychology at the University of North Texas. Current research is the investigation of various aspects of neuropsychological dysfunction in chemically sensitive individuals.

Roy A. Fox, M.D. Environmental Health Center P.O. Box 2130 Fall River, Nova Scotia B2T 1K6 Phone: Fax:

902/860-0057 902/861-1914

Canada

His topics include:

EMF Sensitivity in workers in a hospital with IAO problems and

Heightened Sensitivity Syndrome.

Susan F. Franks

Phone:

817/735-2334

University of North Texas Health Science Center

Fax:

817/735-2299

3500 Camp Bowie Blvd. Department of Psychiatry Ft. Worth, TX 76107

### General Information:

- Dr. Franks received her Ph.D. in Health Psychology & Behavioral Medicine in 1992 from the University of North Texas. She is currently an Assistant Professor in the Dept. of Psychiatry at UNT Health Science Ctr where she is the Director of Neuro-psychological Services and Departmental Director of Research.

Bertie Griffiths, Ph.D.

Phone:

214/368-4132

Environmental Health Center - Dallas

Fax:

214/691-8432

8345 Walnut Hill Lane, Suite 205

Dallas, TX 75231

### General Information:

- Graduate of the University of Wisconsin and the University of the West Indies, Faculty of Medicine.
- Recipient of:
- 1. Degrees in microbiology, virology, and post-doctoral training in Infectious Diseases and Immunology
- 2. Rockefeller fellowship to study Entomology and Virus epidemiology in Brazil a n d Trinidad.
- Appointments:
  - -Professor and Consultant in Microbiology and Infectious Diseases.
  - -Presently Director of EHC-D Analytical Laboratory

Russell Jaffe, M. D., Ph.D. Serammune Physicians Lab 1890 Preston White Dr. Reston VA 20191

Phone:

703/758-0610

Fax:

703/758-0615

General Information: (Education, Specialties, etc.)

- 1972 graduated from Boston University, with a Ph.D. in Biochemistry and his M.D.

- 1972-1973 medical internship University Hospital in Boston

- 1973-1975 residency at Clinical Pathology, National Institutes of Health in Maryland
- He is Board Certified in Pathology and the National Board of Medical Examiners
- 1996 he was awarded Who's Who in Medicine and Healthcare and the POZ 50 Most Innovative AIDS Researchers.

### Additional Information:

-Written a number of books and articles

Wayne Jonas, M.D.
Director of Alternative Medicine
National Institute of Health
9000 Rockville Pike
Bethesda, MD 20892

Phone:

301/594-1990

301/402-2466

Fax:

301/402-4741

General Information: (Education, Specialties, etc.,.)

Dr. Jonas was born in Twin Falls, Idaho. He received his Bachelor of Arts degree from Davidson College in Davidson, NC, and his M.D. from Bowman Gray School of Medicine in Winston-Salem, NC.

Dr. Jonas assumed responsibility of the OAM in July 1995. The OAM was established at the NTH in 1992 to evaluate and investigate complementary, alternative, and unconventional medical practices using appropriate, high quality scientific methods; to coordinate and facilitate research training activities in complementary, alternative, and unconventional medical areas; and to disseminate that information to the public.

Previously, Dr. Jonas served as the Medical Research Fellowship at Walter Reed Army Institute of Research, Washington, D.C.

Alfred R. Johnson, D.O. Environmental Health Center - Dallas 8345 Walnut Hill Lane, Suite 205 Dallas, TX 75231

Phone: 214/368-4132 Fax:

214/691-8432

### General Information:

- Graduate of William Jewell College in Liberty, MO, Received his Doctor of Osteopathic Alfred Medicine at the Univ. Of Health Sciences Center in Kansas City, MO.

- Dr. Johnson has served a six-year term on the Texas State Board of Medical Examiners

- A fellow and former board member of the American Academy of Environmental Medicine, a member of the American College of Osteopathic Internists, the Joint Council of Allergy and Immunology, the Pan American Allergy Society, the American Academy of Disability Evaluating Physicians (fellow), the American Osteopathic Association and the Texas Osteopathic Association. He also holds memberships with American Public Health Association, the American College of Allergists and the American College of Allergy and Immunology.

### Other:

- has published numerous research studies which include topics related to immune system function and examining the health effects of environmental factors on the human body.

Prof. Dr. med. D.G.S. Thilo-Korner Phone: 09 11/97 14-5 31

Euro-Med-Clinic Fax: 11/49-91-19-71-45-32

Europa-Allee 1 90763 Furth

### General Information:

21.08. 1945 Born in Erlangen /FRG

1951-1965 Elementary & High School, Erlangen

1963-1964 Exchange Student, Homer/N.Y./USA

High School Graduate Diploma

1965-1971 University of Erlangen: medicine, psychology

research scholarship from the Faculty of Medicine, University of 1968-1969

Erlangen to the University of Colorado /Denver/USA, Prof. Dr.

K.N. von Kaulla

1970 Finishing medical thesis in Denver 1971 Graduation from University of Erlangen

1972 Physician at I. Medical University Hospital, University of Munich 1974 Married Gabriele

1984 Election to "Hochschulassistent" 1990 Visiting Professor at the University of Rabatt, Могоссо

1994 head of Department of Internal Medicine, Department of Complementary Medicine; Euro-4 - Books, 152 - Scientific Papers, 354 Med-Clinic, Furth /Germany Other:

- Oral Presentations

Rollin McCraty Institute of Heart Math 14700 West Park Avenue Boulder Creek, CA 95006 Phone:

408/338-8727

Fax:

408/338-1182

General Information: (Education, Specialties, etc.,.)

-The Institute of HeartMath is a nonprofit research and educational organization.

-The Institute 's research has laid the foundation for the development of new interventions which enhance human performance and promote physical, mental and emotional well-being.

-Mr. McCraty has an extensive background in systems engineering, radiational physics and instrumentation design which he has brought to the study of human electrophysiology.

-Mr. McCraty has authored a number of scientific papers on heart rate variability, coherent bioelectromagnetic fields and the effects of stress management interventions on human physiology.

Kenneth J. McLeod, Ph.D.

Phone:

516/444-2215 .

State University of New York at Stony Brook

Fax:

516/444-7671

Dept. Orthopedics

H.S.C. T-18 Room 030

Stony Brook, NY 11790-8181

Professor Gavril Markarov

Phone:

095 329 3555

Chief Rehab. Dept.

Fax:

095 395 6377

Hospital 83 Federal Department Med. Biol.

Extreme Problem M. H. Russia

Lipetskaya Str. 13, Apt. 47

Moscow

115598

Russia

General Information: (Education, Specialties, etc.,.)

Twenty years of Magnetolaser treatment of humans in rehabilitation of cardiologies, phlebologies, traumatic, rheumatic, neurologic patients. Twelve years in rehabilitation of radiation exposed patients, EMF Hypersensitivity Neurasthenia Patients by weak EMF, balneotherapy. Seven years of work on bio controlled EMF - chromotherapy of the brain, heart bioelectrical activity, hemodynamics disorder in hypertonic, Parkinson, allergic patients.

1995- present 2nd Professor of Biophys, Dep. Gover. Med. Univer. Moscow

1986- present Chief of Rehab., Physiother. Dep., 83 Hosp., Federal Dep. Med. Bio.

1978-1986 Head of Pysiother. Dep., Physiotherapy Hospital. Moscow